

# RXTE Observations of BL Hya and V2301 Oph

Michael T. Wolff (NRL)

Kent S. Wood (NRL)

James N. Imamura (Univ. of Oregon)

Thomas Steiman-Cameron (Indiana Univ.)

John Middleditch (LANL)

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>1999</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1999 to 00-00-1999</b>	
4. TITLE AND SUBTITLE <b>RXTE Observations of BL Hyi and V2301 Oph</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory, 4555 Overlook Avenue, SW, Washington, DC, 20375</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES <b>13</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# Abstract

We acquired x-ray data for the AM Herculis systems BL Hyi and V2301 Oph using the PCA detector on the Rossi X-Ray Timing Explorer during September-November 1997. We also obtained broad-band optical photometry for BL Hyi using the 1.5m telescope at the Cerro Tololo Inter-American Observatory during September 1997, that was simultaneous with one short stretch of RXTE x-ray observations. In both systems the x-ray emission is consistent with a bright pole and a fainter pole. We find an orbit-modulated optical QPO in BL Hyi consistent with previous results. We characterize the hard x-ray properties of these two systems in this poster.

# BL Hyi Observations

The white dwarf in the BL Hyi system is believed to have a relatively weak magnetic field ( $\mathbf{B}_* \sim 13$  MG; Wickramasinghe et al. 1984) and an orbit period of  $P = 113.7$  minutes (Pickles & Visanathan 1982). During our observations BL Hyi was in an unusual high luminosity state with average visual magnitude  $m_v \sim 14.4$ . We employed the standard FTOOLS package of software to analyze the RXTE data. We have accumulated approximately 50 ksec of RXTE data on this source. The x-ray light curves suggested that two accretion poles were active. The primary x-ray emission region covered  $\sim 45^\circ$  in longitude on the white dwarf. The secondary x-ray emission region flux was 33% of the bright phase flux and was consistent with a point source.


# BL Hyi: Spectra and Light Curve

The x-ray spectra were well-fit by a model consisting of an absorbed bremsstrahlung with Gaussian emission line. For the thermal models, the x-ray spectra had  $kT_x \sim 11.5$  keV, an emission line at  $E \sim 6.80$  keV with equivalent width  $EW = 0.86$  keV, and absorption column  $n_H \sim 1.28 \times 10^{22} \text{ cm}^{-2}$  for the bright pole and  $kT_x \sim 10.8$  keV, an emission line at  $E \sim 6.72$  keV with  $EW \sim 1.25$  keV, and  $n_H \sim 3.23 \times 10^{22} \text{ cm}^{-2}$  for the faint pole. The bright-pole flux was  $2.30 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  over the energy range  $E = 2$  keV to 10 keV and  $7.62 \times 10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$  for the faint pole.

The optical emission showed strong,  $2.7\% \pm 0.2\%$ , quasi-periodic oscillations (QPOs) over the frequency range 0.2 Hz - 0.8 Hz. We found that the QPO amplitude was modulated strongly on the orbital period. The inference that the QPOs were connected with the dominant accretion hot spot was strong. We did not find QPOs in the x-ray emission of BL Hyi. An upper limit of 20% - 25% was placed on 0.2 Hz - 0.8 Hz x-ray QPOs, but we are working to lower this limit. There were no detections of other short period QPOs or coherent features in both the optical and x-ray data.


# BL Hyi Bright Pole Spectrum

Figure 1: Background-corrected PCA hard x-ray spectra for BL Hyi for the bright pole. In addition, the best-fit thermal model and residuals are shown.




# BL Hyi Faint Pole Spectrum

Figure 2: Background-corrected PCA hard x-ray spectra for BL Hyi for the faint pole. In addition, the best-fit thermal model and residuals are shown.



# BL Hyi PCA Light Curve


Figure 3: Background-corrected PCA hard x-ray light curve for BL Hyi. The energy band is 2 keV to 20 keV. The data are binned in intervals of 16 seconds. The orbital phase is based on the ephemeris of Shope and Beuermann (1995).





# BL Hyi: Optical QPO

Figure 4: The time-frequency plot and intensity for our optical observation of BL Hyi on September 24, 1997. The QPO was pulsed  $2.7\% \pm 0.2\%$  over the frequency range 0.2 to 0.8 Hz. Note the strong correlation between QPO amplitude and orbital phase,  $\Phi$ .



# V2301 Oph: Observations

V2301 Oph is an AM Herculis system where the white dwarf has a very weak magnetic field,  $\mathbf{B}_* \sim 7$  MG (Ferrario et al. 1995). The binary period is 113.0 minutes (Siber et al. 1994). The x-ray data were acquired using the PCA detector over the extended period September 27, 1997 to November 24, 1997. We accumulated a total of  $\sim 50$  ksec of data from this source, although that data is scattered over a number of discrete observations due to numerous passages through the South Atlantic Anomaly. We employed the standard package of FTOOLS software to analyze the RXTE data.


# V2301 Oph: Spectra and Light Curve

Figure 5 shows the light curve of V2301 Oph during our RXTE observation of November 22-24, 1997. the light curve shows that our observations caught a number of eclipses of the white dwarf. The system ephemeris used in our analysis is that of Barwig et al. (1994).

The bright pole (pre-eclipse) x-ray spectra were well-fit by an absorbed bremsstrahlung plus Gaussian emission line spectra with  $kT_x \sim 15.2$  keV, an emission line at  $E \sim 6.80$  keV with equivalent width  $EW = 0.38$  keV, and absorption column  $n_H \sim 0.96 \times 10^{22} \text{ cm}^{-2}$ . The x-ray flux was  $3.49 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$  over the energy range  $E = 2$  keV to 10 keV. The faint pole emission, while greater than zero in our data, proved too weak a detection to allow a well constrained fit using XSPEC.


# V2301 Oph PCA Light Curve

Figure 5: Background-corrected PCA hard X-Ray light curve for V2301 Oph. The energy band is 2 keV to 20 keV. The data are binned in intervals of 16 seconds. The orbital phase is based on the ephemeris of Barwig et al. (1994).



# V2301 Oph Bright Pole Spectrum

Figure 6: Background-corrected PCA hard x-ray spectra for V2301 Oph for the bright pole. In addition, the best-fit thermal model and residuals are shown.



# References

- Barwig, H., Ritter, H. & Barnbantner, O. 1994, *A & A*, **288**, 204.
- Ferrario, L., Wickramasinghe, D., Bailey, J. & Bailey, D. 1995, *M.N.R.A.S.*, **273**, 17.
- Pickles, A.J. & Visanathan, N. 1982, *Proc. Astr. Soc. Austr.*, **4**, 425.
- Silber, A.D., Remillard, R.A., Horne, K., & Bradt, H.V. 1994, *Ap. J.*, **424**, 955.
- Matt et al. 1998, *A & A*, **334**, 857.
- Wickramasinghe, D.T. et al. 1984, *Ap. J.*, **286**, 328.